



Denise Wilson & Cassandra Hooper


University of Washington – Seattle
In collaboration with

Coming Alongside -- Environmental Services


National Environmental Health Association
Conference
(NEHA 2012)
June 30, 2012







What is the Mobile Device doing?



Source: The Esoteric Teaching: http://www.oldsite.esotericteaching.org/



What is the Mobile Device emitting?


When the sine wave (carrier + information) gets converted from an electrical signal to a signal "in air", it goes from being an electrical wave to an electromagnetic wave.



The electromagnetic waves are a form of radiation (called radio frequency electromagnetic radiation) that is non-ionizing. Ionizing radiation (x-rays, gamma rays, etc.) is inherently more dangerous than non-ionizing radiation because it is known to directly damage genes/DNA and to lead to uncontrolled reproduction in cells (= cancer). Radio frequency means the carrier (underlying frequency) of the electromagnetic wave is between 100kHz and 10 GHz.



How fast is the Mobile Device radiating?



Typical "carrier" frequencies of mobile devices:

- Cell phones (GSM900) in the U.S.
 - 890-915MHz output & 935-960MHz input
 - Some GSM1900
- WIFI: around 2.4GHz
- Cordless Phone:
 - Older: 1.7 MHz, 43-50 MHz
 - Newer: 900 MHz, 1.9 GHz, 2.4 GHz, 5.8 GHz




How much is the Mobile Device emitting?

Background:

The SAR (Specific Absorption Rate) is the a measure of the rate at which energy is absorbed by the body when exposed to EM radiation. SAR has units of Watts per Kilogram and represents the amount of power absorbed into the whole body or into a small area of tissue (depending on the situation). In cell phones, it makes more sense to express SAR in terms of a small area of tissue (close to the head or in the pants pocket). In the U.S., the FCC says a cell phone should emit no more than 1.6 W/kg over 1 gram of tissue (in contrast, the European Union says no more than 2 W/kg over 10 grams of tissue).


How much is the Mobile Device emitting? Cell Phones:






Other Mobile Devices:

Laptop computers are a particular concern, not because they emit more radiation than base stations, but because they are often used in laps ... in direct contact with the body!








EM Radiation vs. Distance from the Body

Theoretically: basic physics tells us that as the distance between the source of EM radiation and the receiver (the body) increases, the power reaching the body (and resulting radiation absorbed SAR) decreases as the square of that distance.

In Practice: the FCC tests phones at a distance 1.5 cm to 2.5 cm from the body both at the ear and in the pocket. This test does not realistically account for the fact that many people use cell phones "right up to the ear" or stuff phones in their pockets in form fitting or tighter garments than the FCC test presumes.

Implication: the FCC tests may under-estimate the SAR of cell phones in their tests in terms of the way many ordinary consumers use and store their cell phones




EM Radiation vs. Signal Strength:

As the number of "bars" associated with a cell phone call decreases, the signal strength decreases, implying that the cell phone must use more power to transmit the call to the closest cell phone tower. The most common cause of low signal strength is a longer distance to a cell tower, but signal interference (humidity, metal or thick walls) can also reduce signal strength.

Approximate Signal Strength based on "Bars"






Decibels can be easily converted to power. If we assume that the cell phone must directly compensate for loss of signal strength (e.g. a drop in signal strength by a factor of 10 means the phone must transmit 10x the power), then a phone call involving 1 "Bar" can emit up to 3000 times as much power as a phone call at 5 "Bars"

Approximate Signal Strength based on "Bars"







Health Risk #1: Not the Brain

Studies of mice and rats have confirmed that exposure to EM radiation from cell phones damages sperm.

Agarwal et al. Fertility and Sterility 2008


"Use of cell phones decreases the semen quality in men by decreasing the sperm count, motility, viability, and normal morphology."

EM Radiation damages endometrial tissue in female rats through the production of reactive oxygen species.

Gurney et al, Toxicology & Industrial Health 2007

"... oxidative endometrial damage plays an important role in the 900 MHz mobile phone-induced endometrial impairment."

Summary: EM radiation from cell phones stored in pockets near reproductive organs damages reproductive systems for both men and women





Health Risk #2: The Brain

Frei et al, BMJ, 2011

"In this update of a large nationwide cohort study of mobile phone use, there were no increased risks of tumours of the central nervous system, providing little evidence for a causal association."

Khurana et al, Surgical Neurology, 2009

"There is adequate epidemiologic evidence to suggest a link between prolonged cell phone usage and the development of an ipsilateral brain tumor."

Hardell et al, International Journal of Oncology, 2011 "An increased risk was found for
brain tumors> and use of mobile or cordless phone. The risk increased with latency time and cumulative use in hours and was highest in subjects with first use before the age of 20"



Health Risk #2: The Brain

Frei et al, BMJ, 2011

- Pros: Large study population, including those using cell phones over ten years (ten years is an approximate latency for brain tumors/cancer)
- Cons: Study was based on subscriptions rather than actual usage and ignored corporate usage
- Bottom Line: No conclusive link was found, although tumor incidence increased after ten years, especially for women.

Khurana et al, Surgical Neurology, 2009

- Pros: reviews 11 studies, all of which evaluated tumors over periods of greater than ten years and long term cell phone usage; looked at ipsilateral effects only from peer reviewed studies
- Cons: many studies were of smaller populations; selfreport bias of cell phone usage may confound results
- Bottom Line: tumors on the same side of the head as cell phone usage are likely 2X more probable than without heavy cell phone usage.




Health Risk #2: The Brain

Hardell et al, International Journal of Oncology, 2011

- Pros: emphasized those with high cell phone usage (2000 hours over ten years) for greater than ten years, relied on self-report of actual cell phone usage. Studied both ipsilateral and contrlateral tumor incidence.
- Cons: Self-report potentially biased by inaccurate usage reports by side of head; smaller population than Danish study.
- Bottom Line: conclusive link found between ipsilateral brain tumors and cell phone usage

Summary: many cell phone studies are confounded by the relatively "young" state of the technology and "long" latency of brain tumors. Combined with other proven health risks (reproductive harm), however, it is highly likely that evidence linking brain and central nervous system tumors to cell phone usage will continue to increase.





The Bottom Line

Cell Phone Usage can be harmful.

To Reproductive Capacity: YES

To the Brain: PROBABLY

Mobile Devices that communicate with each other wirelessly create a net (likely summative) EM radiation impact on the human body.

Top Five Ways to reduce absorption of EM radiation:

- Make long phone calls over wired phones (either wired land-lines or wired headsets/speaker phone function on cell phones)
- 2. When not in use, store a cell phone at least several inches away from the body (pocket, head, etc.).
- 3. Text if you can. Talk by cell only If you must.
- 4. Avoid long periods close to any base station (wireless router, cell phone tower, etc...).
- 5. Avoid using devices in weak signal locations